

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

**As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.**

Amended by Article 34

WO 98/09948

PCT/EP97/04774

CLAIMS


aph 17
a
a

Use of the following groups of compounds, or their com-

positions, for the preparation of medicaments for the

A method for
 treatment of urinary incontinence *by administering by administering compounds* having the general

1
 formula:




or their salts, where:

A = R(COX)_t where t is an integer 0 or 1;


X = O, NH, NR_{1C} where R_{1C} is a linear or branched alkyl having from 1 to 10 C atoms;

R is chosen from the following groups:

* Group I A), where t = 1,



(IAa)



(IAb)

AMENDED SHEET

where:

R_{III5} is H, a linear or whenever possible branched C_1-C_3 alkyl;

R_{III6} has the same meanings as R_{III5} , or when R_{III5} is H it can be benzyl;

R_{III1} , R_{III2} and R_{III3} are equal or different one from the other and are hydrogen, linear or whenever possible branched C_1-C_6 alkyl or C_1-C_6 alkoxy, or Cl, F, Br;

R_{III4} is R_{III1} or bromine;

preferred are the compounds where R_{III1} , R_{III2} and R_{III4} are H, and R_{III3} is Cl and R_{III3} is in the ortho position to NH; R_{III5} and R_{III6} are H, X is equal to O, and


X_1 is $(CH_2-CH_2-O)_2$;

(I Ab) is the residue of 2-[[2-methyl-3-(trifluoromethyl)phenyl]amino]-3-pyridinecarboxylic acid and when -COOH is present it is known as flunixin.

The compounds preferred are those where $X = O$;

* II A) chosen from the following:

where, when $t = 1$, R is




where R_{2a} and R_{3a} are H, a linear or whenever possible


branched substituted or non-substituted C_1 - C_{12} alkyl,
 allyl, with the proviso that when one of the two is
 allyl the other is H; preferably R_{2a} is H, alkyl has
 from 1 to 4 C atoms, R_{3a} is H;

R_{1a} is chosen from


II Aa)




(II)



(XII)




(IV)




(VII)


*Sub
P1
cont*




(XXXV)




(VI)



(VIII)




(IX)



(X)

668240* 02/24/60



where meanings are as follows:

- in the compounds of formula (IV), residue of ketoprofen:

R_{III1} is H, SR_{III3} where R_{III3} contains from 1 to 4 C linear or whenever possible branched C atoms;

R_{III2} is H, hydroxy;

preferred are the compounds where R_{III1} and R_{III2} are H, R_{3a} is H, and R_{2a} is methyl, $X = O$;

- in the compounds of formula (XXI), residue of carprofen:

R_{XX10} is H, a linear or whenever possible branched alkyl having from 1 to 6 carbon atoms, a C_1 - C_6 alkoxy-carbonyl bound to a C_1 - C_6 alkyl, a C_1 - C_6 carboxyalkyl, a C_1 - C_6 alkanoyl, optionally substituted with halogen, benzyl or halobenzyl, benzoyl or halobenzoyl;

R_{XX11} is H, halogen, hydroxy, CN, a C_1 - C_6 alkyl optionally containing OH groups, a C_1 - C_6 alkoxy, acetyl, benzoyloxy, SR_{XX12} where R_{XX12} is a C_1 - C_6 alkyl; a perfluoroalkyl having from 1-3 C atoms, a C_1 - C_6 carboxyalkyl optio-

Sub
D1
cont

nally containing OH groups, NO₂, sulphamoyl, dialkyl sulphamoyl with the alkyl having from 1 to 6 C atoms, or difluoroalkylsulphonyl with the alkyl having from 1 to 3 C atoms;

R_{xxi1} is halogen, CN, a C₁-C₆ alkyl containing one or more OH groups, a C₁-C₆ alkoxy, acetyl, acetamido, benz-yloxy,

SR_{III3} is as above defined, a perfluoroalkyl having from 1 to 3 C atoms, hydroxy, a carboxyalkyl having from 1 to 6 C atoms, NO₂, amino, a mono- or dialkylamino having from 1 to 6 C atoms, sulphamoyl, a dialkyl sulphamoyl having from 1 to 6 C atoms, or difluoroalkylsulphamoyl as above defined; or R_{xxi} jointly with R_{xxi1} is an alkylene dioxy having from 1 to 6 C atoms;

preferred are the compounds where R_{xxio} is H, the connecting bridge is at position 2, R_{xxi} is H, R_{xxi1} is chlorine and is in the para position to nitrogen;


R_{3a} is H, R_{2a} is methyl and X is O;

- in the compounds of formula (XXXV), residue of thia-profenic acid: Ar is phenyl, hydroxyphenyl optionally mono- or polysubstituted with halogen, an alkanoyl or alkoxy having from 1 to 6 C atoms, a trialalkyl having from 1-6 C atoms, preferably from 1-3 C atoms, cyclo-


- Sol. D, cont*
- pentyl o-hexyl o-heptyl, heterocaryl, preferably thienyl, furyl optionally containing OH, pyridyl;
- the preferred compounds of formula (XXXV) are those where Ar is phenyl, R_{3a} is H, R_{2a} is methyl and X is O;
- in the compounds of formula (II), residue of suprofen, the preferred, where $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$;
 - in the compounds of formula (VI), of which the preferred, indoprofen, when R_{2a} is $\overline{CH_3}$ or indobufen, when R_{2a} is equal to H and $R_{3a} = CH_3$ and $X = O$;
 - in the compounds of formula (VIII), of which the preferred, etodolac, when $R_{3a} = R_{2a} = H$ and $X = O$;
 - in the compounds of formula (VII), of which the preferred, fenoprofen, when $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$;
 - in the compounds of formula (III), of which the preferred, fenbufen, when $R_{3a} = R_{2a} = H$ and $X = O$;
 - in the compounds of formula (X), residue of tolmetin, when $R_{3a} = R_{2a} = H$ and $X = O$;
 - in the compounds of formula (IX), residue of flurbi-

profen, when $R_{3a} = H$, $R_{2a} = CH_3$ and $X = O$;


II Ab):




IIIa)




(XXX)



(XXXI)




(XXXII)




(XXXIII)

Sub
101
cont



(XXXVI)



(XXXVII)

where the meanings are as follows:

- when IIIa) contains $-\text{CH}(\text{CH}_3)-\text{COOH}$ it is known as pranoprofen: α -methyl-5H-[1] benzopyran [2,3-b]pyridine-7-acetic acid; preferred $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$ and $\text{X} = \text{O}$;
- when residue (XXX) contains $-\text{CH}(\text{CH}_3)-\text{COOH}$ it is known as bermoprofen: dibenz [b,f] oxepin-2-acetic acid, preferred is $\text{X} = \text{O}$, $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$;

Sub
D¹
cont

- residue (XXXI) is known as CS-670: 2-[4-(2-oxo-1-cyclohexylidenemethyl)phenyl]propionic acid, when the radical is $-\text{CH}(\text{CH}_3)-\text{COOH}$; preferred $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$ and $\text{X} = \text{O}$;


- residue (XXXII) derives from the known pemedolac which contains group $-\text{CH}_2\text{COOH}$, preferred $\text{R}_{2a} = \text{R}_{3a} = \text{H}$ and $\text{X} = \text{O}$;

- when residue (XXXIII) is saturated with $-\text{CH}_2\text{COOH}$ it is known as pyrazolac: 4-(4-chlorophenyl)-1-(4-fluorophenyl)3-pyrazolyl acid derivatives; preferred $\text{R}_{2a} = \text{R}_{3a} = \text{H}$ and $\text{X} = \text{O}$;

- when residue (XXXVI) is saturated with $-\text{CH}(\text{CH}_3)-\text{COO}-$ it is known as zalcoprofen. When the residue is saturated with a hydroxy or amine group or the acid salts, the compounds are known as dibenzothiepin-derivatives. Preferred $\text{R}_{2a} = \text{H}$, $\text{R}_{3a} = \text{CH}_3$ and $\text{X} = \text{O}$;

- when residue (XXXVII) is CH_2-COOH it derives from the known mofezolac: 3,4-di(p-methoxyphenyl)isoxazol-5-acetic acid; preferred are $\text{R}_{2a} = \text{R}_{3a} = \text{H}$, $t = 1$, $\text{X} = \text{O}$.




* Group IIIA), where $t = 1$,



where:

R_{IVd} and R_{IVd1} are at least one H and the other a linear or whenever possible branched C_1 - C_5 alkyl, preferably C_1 and C_2 , or difluoroalkyl with the alkyl having from 1 to 6 C atoms, preferred is C_1 , or R_{IVd} and R_{IVd1} jointly form a methylene group;

R_{IV} has the following meaning:



where the compounds of group IIIA) have the following meanings:

- in the compounds of formula (II):

*Sal
D
cont*

R_{IV-II} is an alkyl having from 1 to 6 C atoms, a cycloalkyl having from 3 to 7 C atoms, an alcoxymethyl having from 1 to 7 C atoms, a trifluoroalkyl having from 1 to 3 C atoms, vinyl, ethynyl, halogen, an alkoxy having from 1 to 6 C atoms, a difluoroalkoxy with the alkyl having from 1 to 7 C atoms, an alcoxymethyloxy having from 1 to 7 C atoms, an alkylthiomethyloxy with the alkyl having from 1 to 7 C atoms, an alkylmethylthio with the alkyl having from 1 to 7 C atoms, cyano, difluoromethylthio, a substituted phenyl- or phenylalkyl with the alkyl having from 1 to 8 C atoms; preferably R_{IV-II} is CH₃O, R_{IVd} is H and R_{IVd1} is CH₃, and is known as the residue of naproxen;


X = NH and X₁ is equal to (CH₂)₄ or (CH₂CH₂O)₂; also preferred is the same compound where X is equal to O;

- in the preferred compounds of formula (X), for which the residue of loxoprofen has been shown, R_{IVd} is H and R_{IVd1} is CH₃, X = NH or O and X₁ is equal to (CH₂)₄ or (CH₂CH₂O)₂;

- in the compounds of formula (III):


R_{IV-III} is a C₂-C₅ alkyl, even branched when possible, a C₂ and C₃ alkyloxy, allyloxy, phenoxy, phenylthio, a cycloalkyl having from 5 to 7 C atoms, optionally sub-

stituted at position 1 by a C₁-C₂ alkyl;
 preferred is the compound where R_{IV-III} is



and R_{IVd} = H, R_{IVd1} is CH₃, a compound known as the residue of ibuprofen; X = NH and X₁ is equal to (CH₂)₄ or -(CH₂CH₂O)₂; also preferred is the same compound where X = O;

* Group IV A)



(IV)

where A = ROO, t = 1,


of which the residue of the known indomethacin has been shown.

* Group V A) chosen from the following:


- V Aa) fenamates chosen from the following,

where t = 1


*Sub
101
cont*




(V Aa1)



(V Aa2)




(V Aa3)




(V Aa4)

04770-04899


*Sub
P¹
cont*




- V Ab), derivatives of niflumic acid, where $t = 1$:




- V Ac), COX₂ inhibitors, where $t = 0$ and R is as follows:




*Sub
101
cont*



(V Ac2)




(V Ac3)






(V Ac4)


*Sub
D1
cont*



- V Ad) derivatives of diuretics when $t = 1$ and R is as follows:




*Sal
N^o 1
cont*




(V Ad4)

- V Ae) derivatives of diuretics when $t = 0$ and R is as follows:








(V Ae1)




(V Ae2)


Sub
101 cont




*Sub
D¹
cont*




(V Ae8)




(V Ae9)



(V Ae10)



(V Ae11)



(V Ae12)

0914770-042899

*John
10/1
cont*

where the meaning in group V A) is as follows:

- in compounds (V Aa1) the residue of enfenamic acid, 2-[(2-phenylethyl)amino]benzoic acid, has been shown;
- in compounds (V Aa2) the residue of flufenamic acid, 2-[[3-(trifluoromethyl)phenyl]-amino]benzoic acid, has been shown;
- in compounds (V Aa3) the residue of meclofenamic acid, 2-[(2,6-dichloro-3-methylphenyl)amino]benzoic acid, has been shown;
- in compounds (V Aa4) the residue of mefanamic acid, 2-[(2,3-dimethylphenyl)amino]benzoic acid, has been shown;
- in compounds (V Aa5) the residue of tolfenamic acid, 2-[(3-chloro-2-methylphenyl)amino]benzoic acid, has been shown;
- in compounds (V Ab1) the residue of niflumic acid, 2-[[3-(trifluoromethyl)phenyl]amino]-3-pyridine carboxylic acid, has been shown;
- in compounds (V Ac1)_{Rvac1} attached to the oxygen atom in position 2 of the benzene ring of N-(4-nitrophenyl)methansulphonamide can be phenyl or cyclohexane. When R_{vac1} is phenyl the residue is that of nimesulide;
- in compounds (V Ac2) the residue of 3-formylamino-7-

*Sub
101
cont*

methanesulfonylamino-6-phenoxy-4H-1-benzopyran-4-one has been shown;

- in compounds (V Ac3) the atom X_2 that links the radical 2,4-difluorothiophenyl to position 6 of the indanone ring of the residue 5-methanesulfonylamido-1-indanone can be sulfur or oxygen;

- in compounds (V Ac4) the residue of celecoxib 4-[5-(4-methylphenyl)-3-(trifluoromethyl)pyrazol-1-yl] benzenesulphonamide, has been shown;

- in compounds (V Ac5) the residue of 6-[2-(3-ethyl-2,3-dihydro-thiazolyl)thio-5-methanesulphonylamido-3H-isobenzofuran-1-one has been shown.

- in compounds (V Ad1) the residue of bumetanide 3-(Aminosulfonyl)-5-(butylamino)-4-phenoxybenzoic acid has been shown;

- in compounds (V Ad2) the residue of ticrynafen [2,3-Dichloro-4-(2-thienylcarbonyl)-phenoxy]acetic acid has been shown;

- in compounds (V Ad3) the residue of ethacrynic acid [2,3-Dichloro-4-(2-methylene-1-oxobutyl)phenoxy]acetic acid, has been shown;

- in compounds (V Ad4) the residue of piretanide 3-(Aminosulfonyl)-4-phenoxy-5-(1-pyrrolidinyl)benzoic

acid has been shown.

- in compounds (V Ae1) the residue of tripamide (3 α , 4 α , - 7 α , 7 α) -3-(Aminosulphonyl)-4-chloro-N-(octahydro-4,7-metano-2H-isoindol-2-yl)benzamide has been shown.
- in compounds (V Ae2) the residue of torsemide N-[[[(1-Methylethyl)amino]carbonyl]4-[(3-methylphenyl)amino]-3-pyridinesulfonamide has been shown;
- in compounds (V Ae3) the residue of azosemide 2-Chloro-5-(1H-tetrazol-5-yl)-4-[(2-thienylmethyl)amino]benzenesulphonamide has been shown;
- in compounds (V Ae4) the residue of bendroflumethiazide 3,4-Dihydro-3-(phenyl-methyl)-6-(trifluoromethyl)-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has been shown;
- in compounds (V Ae5) the residue of chlorothiazide 6-Chloro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has been shown;
- in compounds (V Ae6) the residue of hydrochlorothiazide 6-Chloro-3,4-dihydro-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has been shown;
- in compounds (V Ae7) the residue of methylclothiazide (6-Chloro-3-(chloromethyl)-3,4-dihydro-2-methyl-2H-1,2,4-benzothiadiazine-7-sulfonamide 1,1-dioxide has

*Sub
D1
cont*


been shown;

- in compounds (V Ae8) the residue of chlorthalidone 2-Chloro-5-(2,3-dihydro-1-hydroxy-3-oxo-1H-isoindol-1-yl)benzensulfonamide has been shown;
- in compounds (V Ae9) the residue of Indapamide 3-(Aminosulfonyl)-4-chloro-N-(2,3-dihydro-2-methyl-1H-indol-1-yl)benzamide has been shown;
- in compounds (VAe10) the residue of metolazone 7-Chloro-1,2,3,4-tetrahydro-2-methyl-3-(2-methylphenyl)-4-oxo-6-quinazolinesulfonamide has been shown;
- in compounds (V Ae11) the residue of quinetazone 7-Chloro-2-ethyl-1,2,3,4-tetrahydro-4-oxo-6-quinazoline-sulfonamide has been shown;
- in compounds (V Ae12) the residue of furosemide 5-(Aminosulfonyl)-4-chloro-2-[(2-furanylmethyl)amino]benzoic acid has been shown.


X_1 in formula A- X_1 -NO₂ is a bivalent connecting bridge chosen from the following:

- YO


where Y is a linear or whenever possible branched C₁-C₂₀ alkylene, preferably having from 2 to 5 carbon atoms, or an optionally substituted cycloalkylene having from 5 to 7 carbon atoms;



where n_3 is an integer from 0 to 3;



where n_4 is an integer from 1 to 6, preferably from 2 to 4;



where R_{1f} = H, CH_3 and n_5 is an integer from 1 to 6, preferably from 2 to 4.

the method
2. ~~Use of the compounds~~ according to Claim 1, in which R is chosen from groups IV A) and V A).

3. Compounds or their compositions for use as medicaments from group V A) in Claim 1.

4. Compounds from group V A) according to Claim 1.

Sub D2
5. Compounds or their compositions for use as medicaments

*Sub
D2
cont*

from group V A) according to Claim 3 for the treatment of musculoskeletal disease of an inflammatory nature, respiratory disease of an inflammatory nature, gynaecological and obstetrical disease including early delivery, pre-eclampsia and dysmenorrhoea, cardiovascular disease including re-stenosis, gastrointestinal tumours.

6. Use of the following compounds, or their compositions, for the preparation of medicaments for the following therapeutical applications:


treatment of respiratory disease: bronchitis, in particular asthma: groups from I A) to V A) in Claim 1;

gynaecological and obstetrical disease including early delivery, pre-eclampsia and dysmenorrhoea: groups from I A) to V A) in Claim 1 and group VI A) as defined below;


vascular disease including re-stenosis: groups from I A) to V A) in Claim 1 and group VI A);

gastrointestinal tumours: groups from I A) to V A) in Claim 1 and group VI A);


the compounds in group VI A) have the general formula



of Claim 1, where $t = 1$, include the following:



(Ia)



(Ib)

where:

R_1 is group $OCOR_3$; where R_3 is methyl, ethyl or a linear or branched C_3-C_5 alkyl, or the residue of a single-ring heterocycle having 5 or 6 atoms which can be aromatic, partially or totally hydrogenated, containing one or more heteroatoms independently chosen from O, N and S; R_2 is hydrogen, hydroxy, halogen, a linear or whenever possible branched alkyl having from 1 to 4 C atoms, a linear or whenever possible branched alcoxyl having from 1 to 4 C atoms; a linear or whenever possible branched perfluoroalkyl having from 1 to 4 C atoms, for example trifluoromethyl, nitro, amino, mono- or

di(C₁₋₄)alkylamino;

R₁ and R₂ jointly are the dioxymethylene group, with the proviso that when X = NH, then X₁ is ethylene and R₂ = H; R₁ cannot be OCOR₃ at position 2 when R₃ is methyl; nI being an integer from 0 to 1;

preferably in Ia), X is equal to O or NH, R₁ is acetoxy, preferably at position 3 or 4, most preferably in the ortho position to CO. X₁ is ethylene or (CH₂CH₂O)₂, R₂ is Hydrogen or halogen, most preferred are the following A X₁ NO₂ compounds: 3-acetoxy-N-(2-nitroxyethyl)-benzamide, 4-acetoxy-N-(2-nitroxyethyl)-benzamide, 3-acetoxy-N-(5-nitroxypenthyl)-benzamide, 2-acetoxy-N-(5-nitroxypenthyl)-benzamide, N-2-(nitroxyethyl)-2-propionoxybenzamide, 2-acetoxy-2-nitroxyethylbenzoate, 2-acetoxy-N-(cis-2-nitroxycyclohexyl)-benzamide, 2-acetoxy-4-chloro-N-(2-nitroxyethyl)-benzamide, N-(2-nitroxyethyl)-2-((4-thiazolindinyl)carbonyloxy)-benzamide hydrochloride, 2-nicotinoyloxy-N-(2-nitroxyethyl)-benzamide, 2-acetoxy-5-nitroxypenthylbenzoate;

preferably in Ib) R₃ = CH₃, nI = 0;

X is equal to O, X₁ is ethylene; in this case Ib) is the residue of acetylsalicylsalicylic acid.

add C₁ / add 91